

Deconfinement phase transition in a two-dimensional model of interacting 2×2 plaquettes

A. Fledderjohann¹, A. Klümper¹ and K.-H. Mütter¹

¹*Physics Department, University of Wuppertal, 42097 Wuppertal, Germany*

A two-dimensional model of interacting plaquettes is studied by means of the real space renormalization group approach. Interactions between the plaquettes are mediated solely by spin excitations on the plaquettes. Depending on the plaquette-plaquette coupling J , we find two regimes:

“confinement” $J_c < J \leq 1$, where the singlet ground state forms an infinite (“confined”) cluster in the thermodynamical limit. Here the singlet-triplet gap vanishes, which is the signature for long range spin-spin correlators.

“deconfinement” $0 \leq J < J_c$, where the singlet ground state “deconfines” - i.e. factorizes - into finite n -clusters of size $2^n \times 2^n$, with $n \leq n_c(J)$. Here the singlet-triplet gap is finite.

The critical value turns out to be $J_c = 0.4822\dots$

PACS numbers: 71.10.Fd, 71.27.+a, 75.10.-b, 75.10.Jm

I. INTRODUCTION

We will discuss in this paper the 2D Hamiltonian

$$H = H_0 + J \cdot H_J \quad (1.1)$$

where H_0 is given by isolated plaquettes occupied with spin-1/2 states and H_J describes nearest neighbour interactions of these plaquettes as shown in Fig. 1. For $J = 1$, the model (1.1) reduces to the well known 2d antiferromagnetic Heisenberg model. The modified Hamiltonian (1.1) has been proposed in studies of structural instabilities of two-dimensional systems.

The singlet-triplet gap (“spin gap”) has been studied by various methods:

- Nonlinear σ model as a low energy effective theory (1), (2)
- modified spin wave theory (3), (4)
- cluster expansion up to fourth order starting from $J = 0$, i.e. isolated plaquettes (5)

In the spin-1/2 case, the model (1.1) is expected (7) to have a quantum phase transition at a critical value J_c^1 , which is signalled by a vanishing singlet-triplet gap for $J > J_c$.

The results of these works are:

- no phase transition for the spin 1/2 case in the nonlinear σ model (8) for any $J \leq 1$.
- $J_c = 0.118$ for the modified spin wave theory [versus $J_c = 0.112$ in linear spin wave theory (6)].
- $J_c = 0.54$ for the cluster expansion.

- A value $J_c = 0.555$ was obtained by means of Ising series expansions (7).
- J_c has been determined by means of the CORE method (contractor renormalization expansion) first by Capponi et al. (9) and recently by Albuquerque et al. (10). These results are somewhat lower $J_c = 0.548$.
- Recent Monte Carlo simulations (11,12) yield values close to $J_c = 0.549$.

The nonlinear σ model approach yields different results for different cut-off schemes. In [Kawakami et al. (8)] no phase transition was found for $S = 1/2$, whereas in [Takano et al. (13)] a critical J_c ($0.2 \leq J_c \leq 0.25$) was obtained.

The phase transition of the magnetic system has been discussed in its correspondence to a superfluid-insulator transition of the boson model (14).

The authors of ref. (10) start from singlet ($S = 0$) and triplet ($S = 1$) plaquette states. Excited states $|S, m\rangle$, $S = 0, 1, 2$, $m = -S, \dots, S$ are absolutely necessary to generate interactions, since singlets alone cannot interact due to total spin conservation.

In a recent paper (15), we have studied how interactions on the 4 plaquette compound - depicted in Fig. 1 - are created by single plaquette excitations. The con-




FIG. 1: 4-plaquette system with $2^n \times 2^n = 4 \times 4$ sites - here $n = 2$; the single plaquette on the right shows the enumeration of plaquette sites.

¹ The critical value J_c is related to the parameter γ_c introduced in (8) by $J_c = (1 - \gamma_c)/(1 + \gamma_c)$

servation of total spin at each interaction point is implemented by means of the Wigner-Eckart Theorem for the transition matrix elements

$$\langle S'_l, m'_l | S_q(x) | S_l, m_l \rangle = v_q \left(\begin{array}{c|c} S'_l & 1 \\ m'_l & q \end{array} \begin{array}{c} S_l \\ m_l \end{array} \right) M(S'_l, x, S_l). \quad (1.2)$$

They can be expressed in terms of a Clebsch-Gordan coefficient and one reduced matrix element $M(S'_l, x, S_l)$. The latter only depends on the initial and final plaquette spin S_l , S'_l and the triplet operator $S_q(x)$ at site x . The phase v_q ($v_+ = -1$, $v_0 = v_- = 1$) results from the transformation properties of the spin operator $S_q(x)$ under the group $SU(2)$. The interaction between neighbouring plaquettes can be traced back to the product of reduced matrix elements at sites x and y (Fig. 2)




FIG. 2: Interaction of neighbouring plaquettes.

$$\overline{M}(S'_l, S_l; S'_r, S_r) = \sum_{\langle x, y \rangle} M(S'_l, x, S_l) M(S'_r, y, S_r) \quad (1.3)$$

Quintuplet excitations ($S = 2$) have not been considered in ref. (10). We want to stress here that triplet-quintuplet transitions are large - comparable with singlet-triplet transitions. It is shown in (15) that the gaps (singlet-triplet and triplet-quintuplet) decrease in the renormalization process. We will see in this paper that the inclusion of quintuplet excitations will move the critical value J_c substantially to a lower value.

The paper is organized as follows:

In Section II we summarize the details of the real space renormalization group approach in 2D models.

In Section III we evaluate the renormalization group flow for various couplings and gaps.

In Section IV we discuss the deconfinement of the ground state wavefunction for $J < J_c$.

In Section V we present a finite size analysis of the singlet-triplet gap in both regimes: $J_c < J < 1$ (confined), $0 \leq J < J_c$ (deconfined).

Section VI is devoted to the staggered magnetization.

II. REAL SPACE RENORMALIZATION GROUP IN 2D MODELS.

In (15) we first studied the interaction matrices $\Delta_S^{(2)}$ of the four plaquette system (Fig. 1) in the sectors with total spin S . The elements of the interaction matrices are fixed on one hand by the Clebsch-Gordan coefficients, which arise in the construction of eigenstates with total

spin S (on the 4-plaquette system) and the evaluation of the Wigner-Eckart Theorem (1.2) for the transition matrix elements. On the other hand $\Delta_S^{(2)}$ only depends on the following couplings

$$\gamma = \frac{1}{a} \cdot \overline{M}(21; 10) \quad (2.1)$$

$$\beta = \frac{1}{a} \cdot \overline{M}(11; 11) \quad (2.2)$$

$$\varepsilon = \frac{1}{a} \cdot \overline{M}(22; 11) \quad (2.3)$$

and gaps

$$\rho = \frac{E_1 - E_0}{a} \quad (2.4)$$

$$\kappa = \frac{E_2 + E_0 - 2E_1}{a} \quad (2.5)$$

We have factored out from the couplings (2.1)-(2.3) the “fundamental” interaction

$$a = \overline{M}(1, 0; 1, 0) \quad (2.6)$$

which is induced by the singlet-triplet transitions on the plaquette.

In Appendix A of ref. (15) one can find the explicit form of the interaction matrices Δ_S (for $J = 1!$) $S = 0, 1, 2$ under the premise that on the four plaquettes only rotational symmetric configurations with singlets, triplets and at most one quintuplet contribute. In this case the dimensions d_S of the interaction matrices Δ_S turn out to be

$$(d_0, d_1, d_2) = (7, 9, 14). \quad (2.7)$$

The factor J in (1.1) is taken into account in the interaction matrices $\Delta_S(\frac{\rho}{J}, \frac{\kappa}{J}, \gamma, \beta, \varepsilon)$ by a rescaling of the normalized gaps (2.4), (2.5) whereas the couplings (2.1)-(2.3) remain unchanged.

Having constructed in this way the interaction matrices $\Delta_S^{(n)}$ on an $n = 2$ cluster ($2^n \times 2^n$) from the ground states on an $n = 1$ cluster, we turned to the question, whether it is possible in general to construct the interaction matrix $\Delta_S^{(n+1)}$, $S = 0, 1, 2$ from the corresponding quantities of a $n \times n$ cluster. This is indeed possible under the assumption that the low energy states on the $(n + 1)$ -cluster can be built up again solely from singlet, triplet, quintuplet ground states on n -clusters. The n -dependence only appears in a renormalization of the couplings (2.1)-(2.3) and energy differences (2.5), (2.5)

Here, we refer to (15) [eqns. (6.1)-(6.4); (6.5), (6.6)] for the used formulas of the renormalization of the couplings and recursion formulas for the scaled energy differences. Note, that J does not appear in the first group of equations, whereas the remaining two (scaled energy differences) are linear in J .

Each step $n \rightarrow n + 1$ in the renormalization procedure demands the diagonalization of the interaction matrices

$\Delta_S^{(n+1)}$, $S = 0, 1, 2$:

$$\Delta_0^{(n+1)}|\sigma^{(n+1)}\rangle = \sigma^{(n+1)}|\sigma^{(n+1)}\rangle \quad (2.8)$$

$$\Delta_1^{(n+1)}|\tau^{(n+1)}\rangle = \tau^{(n+1)}|\tau^{(n+1)}\rangle \quad (2.9)$$

$$\Delta_2^{(n+1)}|\xi^{(n+1)}\rangle = \xi^{(n+1)}|\xi^{(n+1)}\rangle \quad (2.10)$$

The eigenstates $|\sigma^{(n+1)}\rangle$, $|\tau^{(n+1)}\rangle$, $|\xi^{(n+1)}\rangle$ with the largest eigenvalues $\sigma^{(n+1)}$, $\tau^{(n+1)}$, $\xi^{(n+1)}$ enter in the quantities

$$I^{(n+1)}(a, b) \quad , \quad G^{(n+1)}(a, b) \quad (a, b) = (1, 0), (2, 1)$$

$$F_\tau^{(n+1)}(a, a) \quad , \quad F_\xi^{(n+1)}(a, a) \quad (a, a) = (1, 1), (2, 2)$$

according to the bilinear forms:

$$I^{(n+1)}(a, b) = \sum_{k,i} \tau_k^{(n+1)} I_{k,i}(a, b) \sigma_i^{(n+1)} \quad (2.11)$$

$$G^{(n+1)}(a, b) = \sum_{l,k} \xi_l^{(n+1)} G_{l,k}(a, b) \tau_k^{(n+1)} \quad (2.12)$$

$$F_\tau^{(n+1)}(a, a) = \sum_k \left(\tau_k^{(n+1)} \right)^2 F_{\tau,k}(a, a) \quad (2.13)$$

$$F_\xi^{(n+1)}(a, a) = \sum_l \left(\xi_l^{(n+1)} \right)^2 F_{\xi,l}(a, a). \quad (2.14)$$

The contraction $I_{j,i}(1, 0)$, etc. are independent of n and listed in Appendix B of paper (15).

III. NUMERICAL EVALUATION OF THE RENORMALIZATION GROUP FLOW.

We now turn to the numerical evaluation of the recursion formula of the couplings [eqns. (6.1) – (6.4) in (15)] and gaps [eqns. (6.5), (6.6) in (15)] in order to study the n -dependence (i.e. finite size $2^{n+1} \times 2^{n+1}$) and J -dependence. We start with the singlet-triplet gap $\rho^{(n+1)}$, which yields the signature for long range order: From Fig. 3 we see, that there are two different regimes:

a) $J_c \leq J \leq 1$

Here the singlet-triplet gap approaches zero with increasing system size. For $J = 1$ we are close to zero already on small systems for $n_0 = 3$. For decreasing J , $n_0(J)$ increases and seems to diverge for $J \rightarrow J_c$.

b) Below this critical value ($J < J_c$) the singlet-triplet gap $\rho^{(n+1)}$ does not converge to zero anymore. Note also that there is a change in the curvature of $\rho^{(n+1)}$ with n , which is for large n convex if $J_c < J$ but concave if $J_c > J$. This allows for a very precise determination of

$$J_c = 0.4822.. \quad (3.1)$$




FIG. 3: The scaled singlet-triplet gaps $\rho^{(n+1)}$ [(6.5) in (15)] as function of n and J .

Let us next turn to the coupling ratio $\frac{a^{(n+1)}}{2a^{(n)}}$ [(6.1) in (15)]. As function of n this quantity has a maximum, which travels to larger values of n , if J is lowered (Fig. 4).




FIG. 4: The coupling ratio $\frac{a^{(n+1)}}{2a^{(n)}}$ [(6.1) in (15)] as function of n and J .

For $J_c \leq J \leq 1$ all curves approach a common limit for large n :

$$\frac{a^{(n+1)}}{2a^{(n)}} = 0.52. \quad (3.2)$$

For $J \leq J_c$ we observe a monotonic decrease to a limiting value, different from (3.2):

$$\frac{a^{(n+1)}}{2a^{(n)}} = 0.25. \quad (3.3)$$

The n -dependence of the coupling $\gamma^{(n+1)}$ [(6.2) in (15)] is shown in Fig. 5.

For $J_c \leq J \leq 1$ all curves approach a common limit

$$\gamma^{(n+1)} \rightarrow 1.0849.. \quad (3.4)$$




FIG. 5: The coupling $\gamma^{(n+1)}$ [(6.2) in (15)] as function of n and J .

for large n , whereas we observe a monotonic increase with n for $J < J_c$ and a common limit

$$\gamma^{(n+1)} \rightarrow 1.3411.. \quad (3.5)$$

We only want to mention that the “diagonal” couplings [(6.3) and (6.4) in (15)], which do not change the plaquette spins, die out after a few steps.

IV. DECONFINEMENT OF THE GROUND STATE WAVEFUNCTION.

It was pointed out in Section II, that the renormalization group procedure demands in each step $n \rightarrow n+1$ the diagonalization (2.8)-(2.10) of the interaction matrices. We only keep those eigenvectors ($|\sigma^{(n+1)}\rangle$) with largest eigenvalue ($\sigma^{(n+1)}$).

We want to discuss now the physical meaning of the eigenvector components:

$$\sigma_i^{(n+1)} = \langle i, 0; n+1 | \sigma^{(n+1)} \rangle \quad i = 1, \dots, 7 \quad (4.1)$$

in the orthonormal basis $|i, 0; n+1\rangle$ in the singlet sector - defined in Table II of ref. (15). E.g. $(\sigma_1^{(n+1)})^2$ has to be interpreted as the probability to find in the singlet ground state the four plaquette configuration

$$|1, 0\rangle = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \quad (4.2)$$

with four noninteracting singlets.

If

$$(\sigma_1^{(n+1)})^2 \rightarrow 1, \quad (4.3)$$

the singlet ground state $|\sigma^{(n+1)}\rangle$ “deconfines” to the configuration $|1, 0\rangle$ with four noninteracting singlets.

In Fig. 6, we show $(\sigma_1^{(n+1)})^2$ as function of n - i.e. the system size ($2^{n+1} \times 2^{n+1}$).




FIG. 6: The probability $(\sigma_1^{(n+1)})^2$ to find the configuration (4.2) with four noninteracting singlets in the ground state.

For $J_c \leq J \leq 1$, $(\sigma_1^{(n+1)})^2$ converges to

$$(\sigma_1^{(n+1)})^2 \rightarrow 0.110236.. \quad (4.4)$$

For $J < J_c$ the deconfinement limit (4.3) is practically reached at a finite value $n = n_c(J)$, cf. Fig. 7.

$n_c(J)$ decreases with $J < J_c$ and defines the largest cluster size ($2^{n_c(J)} \times 2^{n_c(J)}$) which is still confined.

In summary we can say: In the two-dimensional system of interacting plaquettes defined in (1.1) we observed two phases.

- For $J_c \leq J \leq 1$, there is a confined phase, where the ground state does not factorize into finite n -clusters ($2^n \times 2^n$) but forms one infinite cluster in the thermodynamical limit $n \rightarrow \infty$. The vanishing of the singlet-triplet gap is the characteristic signature of this phase.
- In the deconfined phase $0 \leq J \leq J_c$ the ground state factorizes into finite clusters $n < n_c(J)$ where $n_c(J)$ defines the maximal size of clusters and is shown in Fig. 7.

V. FINITE-SIZE ANALYSIS OF THE SINGLET-TRIPLET GAP.

In the confined regime $J_c \leq J \leq 1$ the singlet-triplet gap

$$E_1^{(n)} - E_0^{(n)} \sim 4^{-n\nu_1} \quad (5.1)$$

vanishes with an exponent

$$\nu_1 = -\frac{\log(1+x)}{\log 4} \quad (5.2)$$




FIG. 7: $n_c(J)$: as a criterion we show for each chosen J -value that number of iterations (n) where $(\sigma_1^{(n+1)})^2$ for the first time exceeds 0.999.

which can be determined from the first derivative

$$x = \frac{d(\tau - \sigma)}{d\rho} = \frac{\partial(\tau - \sigma)}{\partial\rho} + \frac{\partial(\tau - \sigma)}{\partial\kappa} \cdot \frac{d\kappa}{d\rho} \quad (5.3)$$

of the largest eigenvalues σ, τ of the interaction matrices $\Delta_S(\rho, \kappa, \dots)$ in the singlet ($S = 0$) and triplet ($S = 1$) sector:

$$\begin{aligned} \frac{\partial(\tau - \sigma)}{\partial\rho} &= \langle \tau | \frac{\partial\Delta_1}{\partial\rho} | \tau \rangle - \langle \sigma | \frac{\partial\Delta_0}{\partial\rho} | \sigma \rangle \\ &= -4(\tau_1^2 - \sigma_1^2) - 2(\tau_2^2 + \tau_3^2 + \tau_4^2 + \tau_5^2) \\ &\quad + 2(\sigma_2^2 + \sigma_3^2) \end{aligned} \quad (5.4)$$

$$\begin{aligned} \frac{\partial(\tau - \sigma)}{\partial\kappa} &= \langle \tau | \frac{\partial\Delta_1}{\partial\kappa} | \tau \rangle - \langle \sigma | \frac{\partial\Delta_0}{\partial\kappa} | \sigma \rangle \\ &= 1 - \tau_1^2 - \tau_2^2 - \tau_3^2 - \sigma_6^2 - \sigma_7^2. \end{aligned} \quad (5.5)$$

The n - and J -dependence of $\frac{\partial(\tau - \sigma)}{\partial\rho}$ (5.4) - which is the dominant part of (5.3) - is presented in Fig. 8. In the




FIG. 8: The first derivative [(5.3), (5.4)] with respect to ρ .

confined regime $J_c \leq J \leq 1$ all the curves approach a

common limit

$$x^{(n+1)}(J) \rightarrow -\frac{3}{4} \quad (5.6)$$

which leads to a universal exponent (5.2)

$$\nu_1(J) = 1. \quad (5.7)$$

In the deconfined regime $0 \leq J < J_c$ we find a nonvanishing singlet-triplet gap:

$$E_1^{(n)} - E_0^{(n)} = E_1^{(\infty)} - E_0^{(\infty)} + f^{(n)} \quad (5.8)$$

with a finite-size correction

$$f^{(n+1)} - f^{(n)} = J \cdot a^{(n)} (\tau^{(n+1)} - \sigma^{(n+1)}) \quad (5.9)$$

which follows from the difference $\tau^{(n+1)} - \sigma^{(n+1)}$ of the largest eigenvalues $\tau^{(n+1)}, \sigma^{(n+1)}$ in the triplet and singlet sector and the fundamental coupling $a^{(n)}$ (2.6), which can be extracted from Fig. 4. The large n limit of the difference

$$\tau^{(n+1)} - \sigma^{(n+1)} \rightarrow 2 \quad (5.10)$$

turns out to be 2 for all J $0 < J < J_c$ whereas the coupling

$$a^{(n+1)}(J) = \alpha(J) \cdot 2^{-n} \quad (5.11)$$

decreases with the system size $N = 4^n$ as $\frac{1}{\sqrt{N}}$ for all J $0 \leq J < J_c$. $\alpha(J)$ is shown in Fig. 9.




FIG. 9: n -dependence of $\alpha(J) = 2^{-n} \cdot a^{(n+1)}(J)$ - shown after $n = 20$ iterations.

VI. THE STAGGERED MAGNETIZATION.

Finally we want to present our results from the recursion formula [(8.3) in ref. (15)]

$$\begin{aligned}
R^{(n+1)} &= \frac{\langle \sigma^{(n+1)} | \Sigma_-^{(n+1)} \Sigma_+^{(n+1)} | \sigma^{(n+1)} \rangle}{\langle \sigma^{(n)} | \Sigma_-^{(n)} \Sigma_+^{(n)} | \sigma^{(n)} \rangle} \\
&= \sum_{i', i=1}^7 \sigma_{i'}^{(n+1)} \sigma_i^{(n+1)} \Gamma_{i', i}(\gamma^{(n)}) \quad (6.1)
\end{aligned}$$

for the staggered magnetization on an $(n+1)$ -cluster. Note, that the renormalization procedure only enters via the components $\sigma_i^{(n+1)}$, $i = 1, \dots, 7$ on an $(n+1)$ -cluster and the coupling $\gamma^{(n)}$.

The 7×7 matrix $\Gamma_{i', i}(\gamma^{(n)})$ is presented in Appendix C of (15). In Fig. 10 we present the ratio $R^{(n+1)}$ as function of n and J for the case ($d_0 = 7, d_1 = 9, d_2 = 14$) for large n ;




FIG. 10: The ratio $R^{(n+1)}$ as function of n and J for dimensions: $d_0 = 7, d_1 = 9, d_2 = 14$.

all the curves approach a common limit

$$\begin{aligned}
R^{(n)} &\rightarrow 0.7401.. \quad \text{for } J_c \leq J \leq 1 \\
R^{(n)} &\rightarrow 0.25 \quad \text{for } 0 \leq J \leq J_c
\end{aligned} \quad (6.2)$$

VII. DISCUSSION AND PERSPECTIVES.

We have studied in the 2D model with interacting plaquettes various observables like the scaled singlet-triplet gap $\rho^{(n+1)}$ (Fig. 3) as function of n (i.e. system size $2^{n+1} \times 2^{n+1}$) and the coupling parameter J in (1.1). We find spectacular differences in the confinement ($J_c < J \leq 1$) and deconfinement ($J < J_c$) regime, which allows - for a given truncation scheme - for an extremely precise determination of the critical coupling J_c in all these quantities. This means, that the interaction matrices $\Delta_S^{(n+1)}$, $S = 0, 1, 2$ (2.8)-(2.10) and thereby the renormalization group equations [(6.2)-(6.6) in (15)] depend on J in an extremely sensitive way. The reason is a feedback between the scaled energy differences $\rho^{(n)}, \kappa^{(n)}$ - which enter in the diagonals of $\Delta_S^{(n+1)}$ - and the largest eigenvalues $\sigma^{(n+1)}, \tau^{(n+1)}, \xi^{(n+1)}$ (2.8)-(2.10).

This feedback also leads to a dramatic change in the eigenstates $|\sigma^{(n+1)}\rangle, |\tau^{(n+1)}\rangle, |\xi^{(n+1)}\rangle$. E.g. the square of the first component [(4.1) for $i = 1$] $(\sigma_1^{(n+1)})^2$ in the singlet eigenvector $|\sigma^{(n+1)}\rangle$ changes completely if we go from the confined ($J_c \leq J \leq 1$) to the deconfined ($J < J_c$) regime. In the deconfined regime $(\sigma_1^{(n+1)})^2$ is almost one, which means, that the ground state factorizes into 4 noninteracting singlets. In the confined phase $(\sigma_1^{(n+1)})^2$ is very small. Therefore, the remaining components $\sigma_i^{(n+1)}$, $i = 2, \dots, 7$ contribute significantly to the eigenstate $|\sigma^{(n+1)}\rangle$.

These contributions are characterized by excitations of the cluster spins on the four plaquette system. Excitations of cluster spins are necessary to induce cluster-cluster interactions. The vanishing of the singlet-triplet gap - as it is observed in the confinement regime $J_c \leq J \leq 1$ - is a consequence of the cluster-cluster interactions induced by cluster excitations (triplet and quintuplet). We have checked the dependence on the truncation of the interaction matrix by suppressing in Tables II, III, IV (of ref. (15)) all states with one quintuplet plaquette. The dimensions of the interaction matrices reduce to

$$(d_0, d_1, d_2) = (5, 3, 4) \quad (7.1)$$

which of course worsens the renormalization group approach. This is signalled by a somewhat larger singlet-triplet gap. As a consequence the deconfined regime ($J \leq J_c$) is enlarged.

If we look at the deconfinement parameter $(\sigma_1^{(n+1)})^2$, Fig. 11 for the case (7.1), we observe a shift to a larger value of J_c :




FIG. 11: The probability $(\sigma_1^{(n+1)})^2$ evaluated for dimensions (7.1).

$$J_c(5, 3, 4) = 0.5615.. \ ; \ J_c(7, 9, 14) = 0.4822.. \ (7.2)$$

This value is close to the result of ref. (10) obtained without quintuplet excitations. Therefore, the difference in the two values (7.2) reflects the effect of rotational symmetric excited states on the 4 plaquette cluster with one quintuplet. We expect that further excited states

with $n_Q = 2, 3, 4$ quintuplets will lead to changes in the values J_c as well. The Monte Carlo simulations of Janke et al. (12) suggest, that the RG results should converge non-monotonously towards 0.549.

-
- ¹ E. Fradkin, *Field Theory of condensed matter physics*, Addison Wesley (1994); A. M. Tsvelik, *Quantum Field Theory in Condensed Matter Physics*, Cambridge University press, New York (1995)
- ² M. Sénéchal, Phys. Rev. B **47**, 8353 (1993); Phys. Rev. B **48**, 15890 (1993)
- ³ M. Takahashi, Phys. Rev. B **40**, 2494 (1989)
- ⁴ J. E. Hirsch, S. Tang, Phys. Rev. B **40**, 4769 (1989)
- ⁵ R. P. Singh, M. P. Gelfand, A. Huse, Phys. Rev. Lett. **61**, 2484 (1988); H. K. He, C. J. Hamer, J. Oitmaa, J. Phys. A **23**, 1775 (1990); K. Hida, J. Phys. Soc. Jpn. **61**, 1013 (1992)
- ⁶ J. Sirker, A. Klümper, K. Hamacher, Phys. Rev. B **65**, 134409 (2002)
- ⁷ R. P. Singh, Z. Weihong, C. J. Hamer and J. Oitmaa, Phys. Rev. B **60**, 7278 (1999)
- ⁸ A. Koga, S. Kumada, N. Kawakami, J. Phys. Soc. Jpn. **68**, No. 2, 1999; J. Phys. Soc. Jpn. **68**, No. 7, 1999
- ⁹ S. Capponi, A. Läuchli and M. Mambrini, Phys. Rev. B **70**, 1004424 (2004); S. Capponi, Theor. Chem. Acc., **116**, 524 (2006)
- ¹⁰ A. F. Albuquerque, M. Troyer, J. Oitmaa, Phys. Rev. B **78**, 132402 (2008)
- ¹¹ C. H. Aits, U. Löw, A. Klümper, W. Weber, Phys. Rev. B **74**, 014425 (2006)
- ¹² S. Wenzel, L. Bogacz, W. Janke, Phys. Rev. Lett. **101**, 127202 (2008); S. Wenzel, W. Janke, Phys. Rev. B **78**, 064402 (2008), 099902(E) (2008)
- ¹³ K. Takano, Y. Kito, Y. Ōno, and K. Sano, Phys. Rev. Lett. **91**, 197202 (2003)
- ¹⁴ T. Senthil et al., Phys. Rev. B **70**, 144407 (2004)
- ¹⁵ A. Fledderjohann, A. Klümper, K.-H. Mütter, subm. to Eur. Phys. J. (2008)